

## 14C: Vershire-Lombard complex, 8 to 15 percent slopes, rocky

The Vershire, rocky component makes up 50 percent of the map unit. The natural drainage class is well drained. Water movement in the most restrictive layer is low. This component is on hills. The parent material consists of loamy till. Depth to a root restrictive layer, bedrock, lithic, is 20 to 40 inches.

The Lombard, rocky component makes up 35 percent of the map unit. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. This component is on hills. The parent material consists of loamy till over saprolite. Depth to a root restrictive layer, bedrock, lithic, is 60 to 72 inches.

Important farmland classification:	Statewide	Land capability: 3 e	Vermont Agricultural Value Group: 7
------------------------------------	-----------	----------------------	-------------------------------------

## Vermont Residential Onsite Waste Disposal Group and Subgroup: IIc

This unit is moderately suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The depth to bedrock in some areas is the primary concern. A significant percentage of this map unit has sufficient soil depth over bedrock to accept a range of designs. On-site investigations can help avoid areas with limited depth to bedrock. Additional fill material may be needed in some areas in order to meet the separation distance requirement between the bottom of the leachfield and bedrock.

PHYSICAL and CHEMICAL PROPERTIES  EDOSION FACTORS									
0.11	Depth	Depth Typical	Clay	Soil	Permeability (In/Hr)	Organic matter (Pct)	EROSION FACTORS		
Soil name	(In)	texture	(Pct)	reaction (pH)			Kw	Kf	Т
Vershire, rocky	0-8	VFSL	2-15	4.5 - 6.5	0.6-2	2.0-8.0	.43	.43	2
	8-19	VFSL	1-12	4.5 - 6.5	0.6-2	0.5-3.5	.64	.64	
	19-22	VFSL	1-10	4.5 - 7.3	0.6-2	0.5-2.0	.64	.64	
	22-32	UWB			0.01-20				
Lombard, rocky	0-8	VFSL	2-15	5.6 - 7.3	0.6-2	2.0-8.0	.37	.37	5
	8-31	VFSL	1-12	5.6 - 7.3	0.6-2	0.5-3.0	.55	.55	
	31-61	FSL	1-15	6.6 - 7.3	2-6	0.0-0.5	.43	.43	
	61-71	UWB			0.01-20				

		<u>WATE</u>	R FEATURES				SOIL	FEATURES	
	Hydrologic	Depth to seasonal	Flooding		Ponding		Hydric		
Soil name	group	high water table (Feet)	Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)	
Vershire, rocky	С		None		None		No	20-40	
Lombard, rocky	В		None		None		No	60-72	

	LAND USE LIMITA	AGRICULTURAL YIELD DATA				
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre	
Vershire, rocky Lombard, rocky Vershire, rocky Lombard, rocky	Dwellings with basements: Dwellings with basements: Pond reservoir areas: Pond reservoir areas:	,	Depth to hard bedrock Slope Slope Seepage	Pasture Grass-legume hay Corn silage Corn silage Pasture Grass-legume hay	10.5 AUM 4 Tons 18 Tons 18 Tons 10.5 AUM 4 Tons	

	Management	<u>v</u>	VOODLAND MANAGI	<u>EMENT</u>
Soil name	concern	Rating	Reason	Vermont natural communities
/ershire	Harvest equip operability:	Well suited		Northern Hardwood Forest,
ombard	Harvest equip operability:	Well suited		Mesic Red Oak-Northern Hardwood Forest, Rich Northern Hardwood Forest,
/ershire	Road suitability:	Moderately suited	Slope	Hemlock Forest,
ombard	Road suitability:	Moderately suited	Slope	Temperate Acidic Outcrop,
ershire/	Erosion hazard (off-road):	Moderate	Slope/erodibility	Temperate Acidic Cliff, Temperate Calcareous Outcrop,
ombard	Erosion hazard (off-road):	Moderate	Slope/erodibility	Temperate Calcareous Cliff

