

175C: Monadnock fine sandy loam, 0 to 15 percent slopes, extremely bouldery

The Monadnock, extremely bouldery component makes up 79 percent of the map unit. Slopes are 0 to 15 percent. This component is on hills on glaciated uplands, mountains on glaciated uplands. The parent material consists of loamy supraglacial meltout till derived from phyllite and/or granite and gneiss and/or mica schist over sandy and gravelly supraglacial meltout till derived from phyllite and/or granite and gneiss and/or mica schist. Depth to a root restrictive layer, strongly contrasting textural stratification, is 18 to 36 inches (depth from the mineral surface is 17 to 31 inches). The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 85 percent. Below this thin organic horizon the organic matter content is about 2 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria.

Important farmland classification: NPSL

Land capability: 7 s

Vermont Agricultural Value Group: 11

Vermont Residential Onsite Waste Disposal Group and Subgroup: Ic

This unit is well suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. With moderate permeability and slopes less than 20 percent, there are few limitations.

PHYSICAL and CHEMICAL PROPERTIES							EROSION FACTORS		
Soil name	Depth (In)	Typical texture	Clay (Pct)	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	Kw	Kf	T
Monadnock, extremely bouldery	0-3	MPM	---	3.5 - 5.5	1-14	35-95	---	---	3
	3-8	FSL	1-10	3.5 - 6.0	0.1-14	1.0-4.0	.37	.37	
	8-10	FSL	1-10	3.5 - 6.0	0.1-14	2.0-20	.32	.32	
	10-12	FSL	1-10	3.5 - 6.0	0.1-14	2.0-10	.32	.32	
	12-22	GR-FSL	1-10	3.5 - 6.0	0.1-14	1.0-6.0	.24	.43	
	22-25	GR-FSL	1-10	3.5 - 6.0	0.1-14	0.0-2.0	.24	.43	
	25-45	GR-LS	0-1	3.5 - 6.0	1-100	0.0-1.0	.17	.32	
	45-65	GR-LS	0-1	3.5 - 6.0	1-100	0.0-1.0	.17	.32	

WATER FEATURES						SOIL FEATURES		
Soil name	Hydrologic group	Depth to seasonal high water table (Feet)	Flooding		Ponding		Hydric soil?	Depth to bedrock (range in inches)
			Frequency	Duration	Frequency	Duration		
Monadnock, extremely bouldery	B	---	None		None		No	

LAND USE LIMITATIONS				AGRICULTURAL YIELD DATA	
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Monadnock, extremely bouldery	Dwellings with basements:	Not limited			
Monadnock, extremely bouldery	Pond reservoir areas:	Very limited	Seepage		

WOODLAND MANAGEMENT				
Soil name	Management concern	Rating	Reason	Vermont natural communities
Monadnock	Harvest equip operability:	Poorly suited	>3% surface cover fragr	Northern Hardwood Forest, Mesic Red Oak-Northern Hardwood Forest, Beech-Red Maple-Hemlock-Northern Hardwood Forest Variant, Hemlock Forest
Monadnock	Road suitability:	Poorly suited	Rock fragments	
Monadnock	Erosion hazard (off-road):	Slight		