

## CaC: Cabot silt loam, 3 to 15 percent slopes

The Cabot component makes up 82 percent of the map unit. The natural drainage class is poorly drained. Water movement in the most restrictive layer is low. This component is on hills on glaciated uplands, mountains on glaciated uplands. The parent material consists of loamy lodgment till derived from mica schist and/or loamy lodgment till derived from limestone. Depth to a root restrictive layer, densic material, is 6 to 20 inches.

| Important farmland classification: Statewide (b) | Land capability: 3 e | Vermont Agricultural Value Group: 7d |
|--|----------------------|--------------------------------------|
|--|----------------------|--------------------------------------|

## Vermont Residential Onsite Waste Disposal Group and Subgroup: IIId

This unit is marginally suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The depth to the seasonal high water table is the major limitation. A detailed, site-specific analysis is generally required. On-site groundwater level monitoring and determination of induced groundwater mounding is often necessary to establish the suitability of this unit. Curtain drains may help lower the water table to an acceptable level.

| PHYSICAL and CHEMICAL PROPERTIES |       |         |               |                          |                         |                            | EROSION FACTORS |     |   |
|----------------------------------|-------|---------|---------------|--------------------------|-------------------------|----------------------------|-----------------|-----|---|
| Soil name                        | Depth | Typical | Clay<br>(Pct) | Soil<br>reaction<br>(pH) | Permeability<br>(In/Hr) | Organic<br>matter<br>(Pct) | EROSION FACTORS |     |   |
| Soii Hairie                      | (ln)  | texture |               |                          |                         |                            | Kw              | Kf  | Т |
| Cabot                            | 0-7   | SIL     | 1-15          | 5.1 - 7.3                | 0.1-14                  | 3.5-20                     | .43             | .43 | 2 |
|                                  | 7-13  | FSL     | 1-15          | 5.1 - 7.3                | 0.1-14                  | 0.3-3.5                    | .49             | .49 |   |
|                                  | 13-65 | FSL     | 1-15          | 5.6 - 7.8                | 0.001-1                 | 0.2-1.0                    | .43             | .43 |   |

| WATER FEATURES |            |                            |           |          | SOIL FEATURES |          |        |                                    |
|----------------|------------|----------------------------|-----------|----------|---------------|----------|--------|------------------------------------|
|                | Hydrologic | Depth to seasonal          |           |          | Ponding       |          | Hydric |                                    |
| Soil name      | group      | high water table<br>(Feet) | Frequency | Duration | Frequency     | Duration | soil?  | Depth to bedrock (range in inches) |
| Cabot          | D          | 0.0-1.5                    | None      |          | None          |          | Yes    |                                    |

|                        | LAND USE LIMITA           | AGRICULTURAL YIELD DATA |                         |                  |              |
|------------------------|---------------------------|-------------------------|-------------------------|------------------|--------------|
| Soil name              | Land use                  | Rating                  | Reason **               | Crop name        | Yield / acre |
| Cabot                  | Dwellings with basements: | Very limited            | Depth to saturated zone | Grass-legume hay | 3.5 Tons     |
| Cabot                  | Pond reservoir areas:     | Very limited            | Slope                   | Grass-clover     | 5.6 AUM      |
| Total received direct. |                           | vory miniod             | Clopo                   | Corn silage      | 13 Tons      |

|           | Management                 | WOODLAND MANAGEMENT |                                 |   |  |  |
|-----------|----------------------------|---------------------|---------------------------------|---|--|--|
| Soil name | concern                    | Rating              | Reason                          | Vermont natural communities                                     |  |  |
| Cabot     | Harvest equip operability: | Poorly suited       | <30cm to water table for >=6mos | Red Maple-Black Ash Swamp,<br>Spruce-Fir-Tamarack Swamp,        |  |  |
| Cabot     | Road suitability:          | Moderately suited   | Slope                           | Lowland Spruce-Fir Forest, Alder Swamp,                         |  |  |
| Cabot     | Erosion hazard (off-road): | Slight              |                                 | Northern White Cedar Swamp, Calcareous Red Maple-Tamarack Swamp |  |  |