

MvB: Munson and Raynham silt loams, 2 to 6 percent slopes

The Munson component makes up 45 percent of the map unit. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is very low. This component is on terraces on lake plains. The parent material consists of coarse-silty glaciolacustrine deposits over clayey glaciolacustrine deposits. Depth to a root restrictive layer is greater than 60 inches

The Raynham component makes up 45 percent of the map unit. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. This component is on terraces on lake plains. The parent material consists of coarse-silty glaciolacustrine deposits. Depth to a root restrictive layer is greater than 60 inches.

Important farmland classification: Statewide (b)	Land capability: 3 w	Vermont Agricultural Value Group: 4d
--	----------------------	--------------------------------------

Vermont Residential Onsite Waste Disposal Group and Subgroup: IVa

This unit is generally not suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. Excessive soil wetness in association with the minimal slope is the limiting condition. Prolonged periods of saturation at or near the soil surface do not allow for the proper functioning of septic systems.

PHYSICAL and CHEMICAL PROPERTIES EROSION E							ION EA	CTOPS	
Cailmana	Depth	Typical	Clay	Soil	Permeability	Organic matter	EROSION FACTORS		
Soil name	(In) texture (Pct) reaction (pH)		(In/Hr)	Kw	Kf	Т			
Munson	0-8	SIL	3-10	5.6 - 6.5	0.6-2	3.0-10	.49	.49	3
	8-15	SIL	3-16	5.6 - 6.5	0.2-2	0.5-3.0	.64	.64	
	15-65	SIC	35-60	5.6 - 7.3	0-0.2	0.0-1.0	.32	.32	
Raynham	0-6	SIL	3-16	5.1 - 7.3	0.2-2	3.0-10	.37	.37	5
	6-22	SIL	3-16	5.1 - 7.3	0.2-2	0.5-2.0	.64	.64	
	22-65	SIL	3-16	5.6 - 7.8	0.06-0.2	0.0-0.5	.64	.64	

		WATE	R FEATURES				SOIL	<u>FEATURES</u>	
	Hydrologic	Depth to seasonal	Flooding		Ponding		Hydric		
Soil name	group	high water table (Feet)	Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)	
Munson	C/D	0.5-2.0	None		None		No		
Raynham	C/D	0.0-2.0	None		None		Yes		

	LAND USE LIMITA	TIONS		AGRICULTURAL YIE	LD DATA
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Munson	Dwellings with basements:	Very limited	Depth to saturated zone	Corn silage	18 Tons
Raynham	Dwellings with basements:	Very limited	Depth to saturated zone	Grass hay	4 Tons
Munson	Pond reservoir areas:	Somewhat limited	Slope	Grass-clover	5.6 AUM
Raynham		Somewhat limited	•	Grass-legume hay	3.5 Tons
Kayıllallı	Pond reservoir areas:	Somewhat iimited	Seepage	Grass-legume hay	3.5 Tons
				Grass-clover	5.6 AUM
				Grass hay	4 Tons
				Corn silage	22 Tons

	Management	<u>v</u>	<u>VT</u>	
Soil name	concern	Rating	Reason	Vermont natural communities
Munson	Harvest equip operability:	Poorly suited	<30cm to water table for >=6mos	Valley Clayplain Forest
Raynham	Harvest equip operability:	Moderately suited	30-60cm to water table t	
Munson	Road suitability:	Poorly suited	Wetness	
Raynham	Road suitability:	Moderately suited	Wetness	

Soil Fact Sheet - Continued

Chittenden County, Vermont

Munson Erosion hazard (off-road): Slight Raynham Erosion hazard (off-road): Slight