

**MyC: Munson and Raynham silt loams, 6 to 12 percent slopes**

The Munson component makes up 45 percent of the map unit. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is very low. This component is on terraces on lake plains. The parent material consists of coarse-silty glaciolacustrine deposits over clayey glaciolacustrine deposits. Depth to a root restrictive layer is greater than 60 inches.

The Raynham component makes up 45 percent of the map unit. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. This component is on terraces on lake plains. The parent material consists of coarse-silty glaciolacustrine deposits. Depth to a root restrictive layer is greater than 60 inches.

Important farmland classification: Statewide (b)	Land capability: 3 e	Vermont Agricultural Value Group: 7d
--	----------------------	--------------------------------------

Vermont Residential Onsite Waste Disposal Group and Subgroup: III d

This unit is marginally suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The depth to the seasonal high water table is the major limitation. A detailed, site-specific analysis is generally required. On-site groundwater level monitoring and determination of induced groundwater mounding is often necessary to establish the suitability of this unit. Curtain drains may help lower the water table to an acceptable level.

PHYSICAL and CHEMICAL PROPERTIES							EROSION FACTORS		
Soil name	Depth (In)	Typical texture	Clay (Pct)	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	Kw	Kf	T
Munson	0-8	SIL	3-10	5.6 - 6.5	0.6-2	3.0-10	.49	.49	3
	8-15	SIL	3-16	5.6 - 6.5	0.2-2	0.5-3.0	.64	.64	
	15-65	SIC	35-60	5.6 - 7.3	0-0.2	0.0-1.0	.32	.32	
Raynham	0-6	SIL	3-16	5.1 - 7.3	0.2-2	3.0-10	.37	.37	5
	6-22	SIL	3-16	5.1 - 7.3	0.2-2	0.5-2.0	.64	.64	
	22-65	SIL	3-16	5.6 - 7.8	0.06-0.2	0.0-0.5	.64	.64	

WATER FEATURES						SOIL FEATURES		
Soil name	Hydrologic group	Depth to seasonal high water table (Feet)	Flooding		Ponding		Hydric soil?	Depth to bedrock (range in inches)
			Frequency	Duration	Frequency	Duration		
Munson	C/D	0.5-2.0	None		None		No	---
Raynham	C/D	0.0-2.0	None		None		Yes	---

LAND USE LIMITATIONS				AGRICULTURAL YIELD DATA	
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Munson	Dwellings with basements:	Very limited	Depth to saturated zone	Grass-legume hay	3.5 Tons
Raynham	Dwellings with basements:	Very limited	Depth to saturated zone	Grass-clover	5.6 AUM
Munson	Pond reservoir areas:	Very limited	Slope	Grass hay	4 Tons
Raynham	Pond reservoir areas:	Very limited	Slope	Corn silage	18 Tons
				Corn silage	20 Tons
				Grass-legume hay	3.5 Tons
				Grass-clover	5.6 AUM
				Grass hay	4 Tons

WOODLAND MANAGEMENT				
Soil name	Management concern	Rating	Reason	Vermont natural communities
Munson	Harvest equip operability:	Poorly suited	<30cm to water table for >=6mos	Valley Clayplain Forest
Raynham	Harvest equip operability:	Moderately suited	30-60cm to water table for >=6mos	
Munson	Road suitability:	Poorly suited	Wetness	
Raynham	Road suitability:	Moderately suited	Wetness	

---

Munson	Erosion hazard (off-road):	Slight
Raynham	Erosion hazard (off-road):	Slight