

Br: Birdsall silt loam

The Birdsall component makes up 85 percent of the map unit. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. This component is on depressions on lake plains. The parent material consists of coarse-silty glaciolacustrine deposits. Depth to a root restrictive layer is greater than 60 inches.

Important farmland classification: NPSL	Land capability: 5 w	Vermont Agricultural Value Group: 10
---	----------------------	--------------------------------------

Vermont Residential Onsite Waste Disposal Group and Subgroup: IVa

This unit is generally not suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. Excessive soil wetness in association with the minimal slope is the limiting condition. Prolonged periods of saturation at or near the soil surface do not allow for the proper functioning of septic systems.

PHYSICAL and CHEMICAL PROPERTIES								EDOSION EACTORS	
Depth Typical texture	Clay (Pct)	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	EROSION FACTORS				
					Kw	Kf	Т		
0-5	SIL	3-16	4.5 - 6.0	0.2-2	2.0-8.0	.49	.49	5	
5-21	VFSL	3-16	5.1 - 7.3	0.2-0.6	0.5-3.0	.55	.55		
21-60	SIL	3-16	5.1 - 7.3	0.06-0.2	0.0-1.0	.64	.64		
	0-5 5-21	Depth (In) Typical texture 0-5 SIL 5-21 VFSL	Depth (In) Typical texture Clay (Pct) 0-5 SIL 3-16 5-21 VFSL 3-16	Depth (In) Typical texture Clay (Pct) Soil reaction (pH) 0-5 SIL 3-16 4.5 - 6.0 5-21 VFSL 3-16 5.1 - 7.3	Depth (In) Typical texture Clay (Pct) Soil reaction (pH) Permeability (In/Hr) 0-5 SIL 3-16 4.5 - 6.0 0.2-2 5-21 VFSL 3-16 5.1 - 7.3 0.2-0.6	Depth (In) Typical texture Clay (Pct) Soil reaction (pH) Permeability (In/Hr) Organic matter (Pct) 0-5 SIL 3-16 4.5 - 6.0 0.2-2 2.0-8.0 5-21 VFSL 3-16 5.1 - 7.3 0.2-0.6 0.5-3.0	Depth (In) Typical texture Clay (Pct) Soil reaction (pH) Permeability (In/Hr) Organic matter (Pct) Kw 0-5 SIL 3-16 4.5 - 6.0 0.2-2 2.0-8.0 .49 5-21 VFSL 3-16 5.1 - 7.3 0.2-0.6 0.5-3.0 .55	Depth (In) Typical texture Clay (Pct) Soil reaction (pH) Permeability (In/Hr) Organic matter (Pct) Kw Kf 0-5 SIL 3-16 4.5 - 6.0 0.2-2 2.0-8.0 .49 .49 5-21 VFSL 3-16 5.1 - 7.3 0.2-0.6 0.5-3.0 .55 .55	

WATER FEATURES							SOIL FEATURES	
Soil name	Hydrologic group	Depth to seasonal high water table (Feet)	Flooding		Ponding		Hydric	
			Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)
Birdsall	C/D	0.0-0.5	None		Frequent	Long (7 to 30 days)	Yes	

	LAND USE LIMITATION	AGRICULTURA	L YIELD DATA			
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre	
Birdsall	Dwellings with hasements: Ve	ny limited	Ponding			

Birdsall Dwellings with basements: Very limited Ponding

Birdsall Pond reservoir areas: Somewhat limited Seepage

	Management		<u>NT</u>	
Soil name	concern	Rating	Reason	Vermont natural communities
Birdsall	Harvest equip operability:	Poorly suited	<30cm to water table fo >=6mos	Red Maple-Black Ash Swamp, Alder Swamp,Sedge Meadow
Birdsall	Road suitability:	Poorly suited	Ponding	
Birdsall	Erosion hazard (off-road):	Slight		•