

HbB: Hinesburg loamy fine sand, 3 to 8 percent slopes

The Hinesburg component makes up 65 percent of the map unit. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. This component is on terraces on lake plains. The parent material consists of sandy glaciofluvial deposits over loamy glaciolacustrine deposits. Depth to a root restrictive layer is greater than 60 inches.

Important farmland classification: Prime	Land capability: 2 s	Vermont Agricultural Value Group: 3
--	----------------------	-------------------------------------

Vermont Residential Onsite Waste Disposal Group and Subgroup: Ilh

This unit is moderately suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The depth to the seasonal high water table is the primary concern. Mound system construction and other site modifications are often necessary. On sloping sites, curtain drains can help lower the water table to an acceptable level. In some cases, a detailed, site-specific analysis with groundwater level monitoring and determination of induced groundwater mounding may be required to establish the suitability of this unit.

PHYSICAL and CHEMICAL PROPERTIES							EROSION FACTORS		
Soil name	Depth Typ	Typical	Clay (Pct)	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	EROSION FACTORS		
	(ln)						Kw	Kf	Т
Hinesburg	0-7	LFS	1-5	5.6 - 6.5	6-20	3.0-6.0	.17	.17	5
	7-22	LFS	1-5	5.6 - 6.5	6-20	0.5-2.0	.37	.37	
	22-60	SIL	3-28	5.1 - 7.3	0.2-0.6	0.0-0.5	.64	.64	

WATER FEATURES						SOIL FEATURES		
	Hydrologic Depth to seasonal		Floo	Flooding		Ponding		
Soil name	group	high water table (Feet)	Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)
Hinesburg	Α	2.0-4.0	None		None		No	

	LAND USE LIMITA	AGRICULTURAL YIELD DATA			
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Hinesburg	Dwellings with basements:	Somewhat limited	Depth to saturated zone	Alfalfa hay	4 Tons
Hinesburg	Pond reservoir areas:	Very limited	Seepage	Grass-legume hay	3.5 Tons
riiiosburg	Fond reservoir areas.	very innited	Occpage	Grass-clover	5.6 AUM
				Grass hay	3 Tons
				Corn silage	16 Tons

	Management		WOODLAND MANA	AGEMENT
Soil name	concern	Rating	Reason	Vermont natural communities
Hinesburg	Harvest equip operability:	Well suited		White Pine-Red Oak-Black Oak Forest,
Hinesburg	Road suitability:	Well suited		White Pine-Northern Hardwood Forest Variant, Sugar Maple-Ostrich Fern Riverine Floodplain
Hinesburg	Erosion hazard (off-road):	Slight		Forest