

NnB: Ninigret fine sandy loam, 0 to 8 percent slopes

The Ninigret component makes up 85 percent of the map unit. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. This component is on depressions on terraces on river valleys. The parent material consists of coarse-loamy glaciofluvial deposits over sandy and gravelly glaciofluvial deposits. Depth to a root restrictive layer is greater than 60 inches.

Important farmland classification: Prime

Land capability: 2 w

Vermont Agricultural Value Group: 3

Vermont Residential Onsite Waste Disposal Group and Subgroup: IIh

This unit is moderately suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The depth to the seasonal high water table is the primary concern. Mound system construction and other site modifications are often necessary. On sloping sites, curtain drains can help lower the water table to an acceptable level. In some cases, a detailed, site-specific analysis with groundwater level monitoring and determination of induced groundwater mounding may be required to establish the suitability of this unit.

PHYSICAL and CHEMICAL PROPERTIES							EROSION FACTORS		
Soil name	Depth (In)	Typical texture	Clay (Pct)	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	Kw	Kf	T
Ninigret, frigid	0-8	FSL	2-8	4.5 - 6.0	0.1-14	2.0-5.0	.32	.32	3
	8-16	FSL	2-8	4.5 - 6.0	0.1-14	0.5-1.5	.37	.37	
	16-26	FSL	2-8	4.5 - 6.0	0.1-14	0.0-0.5	.43	.43	
	26-65	SR- LS LFS	1-5	4.5 - 6.5	1-100	0.0-0.5	.43	.43	

WATER FEATURES						SOIL FEATURES		
Soil name	Hydrologic group	Depth to seasonal high water table (Feet)	Flooding		Ponding		Hydric soil?	Depth to bedrock (range in inches)
			Frequency	Duration	Frequency	Duration		
Ninigret, frigid	C	1.4-3.2	None		None		No	---

LAND USE LIMITATIONS				AGRICULTURAL YIELD DATA	
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Ninigret, frigid	Dwellings with basements:	Very limited	Depth to saturated zone	Corn silage	30 Tons
				Pasture	5 AUM
				Tobacco	1850 Lbs
				Grass hay	3.5 Tons
				Corn	158 Bu
				Grass-legume hay	3.5 Tons

WOODLAND MANAGEMENT				
Soil name	Management concern	Rating	Reason	Vermont natural communities
Ninigret	Harvest equip operability:	Well suited		White Pine-Red Oak-Black Oak Forest, White Pine-Northern Hardwood Forest Variant, Sugar Maple-Ostrich Fern Riverine Floodplain Forest
Ninigret	Road suitability:	Well suited		
Ninigret	Erosion hazard (off-road):	Slight		