

393D: Monadnock fine sandy loam, 15 to 35 percent slopes, extremely bouldery

The Monadnock, extremely bouldery component makes up 80 percent of the map unit. Slopes are 15 to 35 percent. This component is on hills on glaciated uplands, mountains on glaciated uplands. The parent material consists of loamy supraglacial meltout till derived from phyllite and/or granite and gneiss and/or mica schist over sandy and gravelly supraglacial meltout till derived from phyllite and/or granite and gneiss and/or mica schist. Depth to a root restrictive layer, strongly contrasting textural stratification, is 18 to 36 inches (depth from the mineral surface is 17 to 31 inches). The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 85 percent. Below this thin organic horizon the organic matter content is about 2 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria.

Important farmland classification: NPSL

Land capability: 7 s

Vermont Agricultural Value Group: 11

Vermont Residential Onsite Waste Disposal Group and Subgroup: Id

This unit is well suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. Slopes greater than 20 percent in some areas are a concern. There may be less-sloping areas within the unit that are suitable for siting a septic system, or, if feasible, cut and fill site modifications may produce an acceptable area within the unit. An erosion prevention and sediment control plan is required by the State for construction on sites over 20 percent slope.

PHYSICAL and CHEMICAL PROPERTIES							EROSION FACTORS		
Soil name	Depth (In)	Typical texture	Clay (Pct)	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	Kw	Kf	T
Monadnock, extremely bouldery	0-3	MPM	---	3.5 - 5.5	1-14	35-95	---	---	3
	3-8	FSL	1-10	3.5 - 6.0	0.1-14	1.0-4.0	.37	.37	
	8-10	FSL	1-10	3.5 - 6.0	0.1-14	2.0-20	.32	.32	
	10-12	FSL	1-10	3.5 - 6.0	0.1-14	2.0-10	.32	.32	
	12-22	GR-FSL	1-10	3.5 - 6.0	0.1-14	1.0-6.0	.24	.43	
	22-25	GR-FSL	1-10	3.5 - 6.0	0.1-14	0.0-2.0	.24	.43	
	25-45	GR-LS	0-1	3.5 - 6.0	1-100	0.0-1.0	.17	.32	
	45-65	GR-LS	0-1	3.5 - 6.0	1-100	0.0-1.0	.17	.32	

WATER FEATURES						SOIL FEATURES	
Soil name	Hydrologic group	Depth to seasonal high water table (Feet)	Flooding		Ponding		Hydric soil?
			Frequency	Duration	Frequency	Duration	
Monadnock, extremely bouldery	B	---	None		None		No

LAND USE LIMITATIONS				AGRICULTURAL YIELD DATA	
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Monadnock, extremely bouldery	Dwellings with basements:	Very limited	Slope		
Monadnock, extremely bouldery	Pond reservoir areas:	Very limited	Slope		

WOODLAND MANAGEMENT				
Soil name	Management concern	Rating	Reason	Vermont natural communities
Monadnock	Harvest equip operability:	Poorly suited	>3% surface cover fragr	Northern Hardwood Forest,
Monadnock	Road suitability:	Poorly suited	Slope	Mesic Red Oak-Northern Hardwood Forest,
Monadnock	Erosion hazard (off-road):	Moderate	Slope/erodibility	Beech-Red Maple-Hemlock-Northern Hardwood Forest Variant, Hemlock Forest