

58C: Colton-Duxbury complex, 8 to 15 percent slopes

The Colton component makes up 50 percent of the map unit. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. This component is on knolls on river valleys, terraces on river valleys. The parent material consists of sandy and gravelly glaciofluvial deposits. Depth to a root restrictive layer is greater than 60 inches.

The Duxbury component makes up 35 percent of the map unit. The natural drainage class is well drained. Water movement in the most restrictive layer is high. This component is on knolls on river valleys, terraces. The parent material consists of coarse-loamy glaciofluvial deposits over sandy and gravelly glaciofluvial deposits. Depth to a root restrictive layer is greater than 60 inches.

Important farmland classification: NPSL	Land capability: 4 e	Vermont Agricultural Value Group: 8
---	----------------------	-------------------------------------

Vermont Residential Onsite Waste Disposal Group and Subgroup: la

This unit is well suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The rapid permeability in the substratum is a concern. Backfilling absorption trenches with at least one foot of finer textured material or other site modifications may be necessary to slow the percolation rate enough to allow for thorough filtering of effluent.

PHYSICAL and CHEMICAL PROPERTIES							EROSION FACTORS		
Cailnama	Depth	Typical	Clay	Soil	Permeability	Organic	<u>EROSION FACTORS</u>		
Son name	Soil name Deptit Typical Clay reaction (Pct) (pH)		(In/Hr) matter (Pct)		Kw	Kf	Т		
Colton	0-6	GRV-SL	1-7	4.5 - 6.0	6-20	2.0-6.0	.05	.17	5
	6-32	GRX-S	0-5	4.5 - 6.0	6-20	0.5-2.0	.02	.02	
	32-60	GRX-S	0-3	4.5 - 6.5	20-100	0.0-0.5	.02	.02	
Duxbury	0-5	FSL	1-7	4.5 - 6.5	2-6	2.0-5.0	.24	.24	2
	5-18	GR-FSL	2-8	4.5 - 6.5	2-6	0.5-3.0	.24	.37	
	18-60	GRV-LS	0-2	4.5 - 6.5	6-20	0.0-1.0	.10	.24	

		<u>WATE</u>	R FEATURES				SOIL	<u>FEATURES</u>
	Hydrologic	Depth to seasonal	Floo	ding	Pon	ding	Hydric	
Soil name	group	high water table (Feet)	Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)
Colton	Α		None		None		No	
Duxbury	Α		None		None		No	

	LAND USE LIMITAT	AGRICULTURAL YI	AGRICULTURAL YIELD DATA			
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre	
Colton	Dwellings with basements:	Somewhat limited	Slope	Grass-legume hay	2 Tons	
Duxbury	Dwellings with basements:	Somewhat limited	Slope	Alfalfa hay	2.5 Tons	
Colton	G	Very limited	Seepage	Grass-clover	3.2 AUM	
	Pond reservoir areas:	,	. •	Grass hay	2 Tons	
Duxbury	Pond reservoir areas:	Very limited	Seepage	Grass-clover	4 AUM	
				Grass hay	2.5 Tons	
				Corn silage	18 Tons	
				Alfalfa hay	3 Tons	

	Management	<u>w</u>	OODLAND MANA	AGEMENT
Soil name concern	· ·	Rating	Reason	Vermont natural communities
Colton	Harvest equip operability:	Well suited		Hemlock-Northern Hardwood Forest,
Duxbury	Harvest equip operability:	Well suited		Hemlock-White Pine-Northern Hardwood Forest Variant.
Colton	Road suitability:	Moderately suited	Slope	Lowland Spruce-Fir Forest, well drained phase,
Duxbury	Road suitability:	Moderately suited	Slope	White Pine-Northern Hardwood Forest Variant,
Colton	Erosion hazard (off-road):	Slight		Hemlock Forest, Red Pine Forest or Woodland
Duxbury	Erosion hazard (off-road):	Slight		

