

66C: Georgia and Amenia soils, 8 to 15 percent slopes

The Amenia component makes up 40 percent of the map unit. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately low. This component is on hills on glaciated uplands, knolls on glaciated uplands, ridges on glaciated uplands. The parent material consists of coarse-loamy till. Depth to a root restrictive layer is greater than 60 inches.

The Georgia component makes up 40 percent of the map unit. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately low. This component is on hills on glaciated uplands, knolls on glaciated uplands, ridges on glaciated uplands. The parent material consists of coarse-loamy till. Depth to a root restrictive layer is greater than 60 inches.

Important farmland classification:	Statewide	Land capability: 3 e	Vermont Agricultural Value Group: 7
------------------------------------	-----------	----------------------	-------------------------------------

Vermont Residential Onsite Waste Disposal Group and Subgroup: IIh

This unit is moderately suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The depth to the seasonal high water table is the primary concern. Mound system construction and other site modifications are often necessary. On sloping sites, curtain drains can help lower the water table to an acceptable level. In some cases, a detailed, site-specific analysis with groundwater level monitoring and determination of induced groundwater mounding may be required to establish the suitability of this unit.

PHYSICAL and CHEMICAL PROPERTIES							EROSION FACTORS		
Soil name	Depth	Typical	Clay	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	<u>EROSION FACTORS</u>		
	(In)	. , , , , , , , , , , , , , , , , , , ,	(Pct)				Kw	Kf	Т
Amenia	0-8	L	5-18	5.6 - 7.3	0.6-2	2.0-6.0	.28	.28	3
	8-24	GR-L	5-18	5.6 - 7.3	0.6-2	0.5-2.0	.28	.49	
	24-60	GR-FSL	5-18	7.4 - 8.4	0.06-0.2	0.0-0.5	.24	.37	
Georgia	0-9	L	7-18	5.1 - 7.3	0.6-2	3.0-8.0	.28	.28	3
	9-30	GR-L	5-18	5.1 - 7.3	0.6-2	0.0-1.0	.28	.43	
	30-60	GR-L	10-18	5.1 - 7.3	0.06-0.2	0.0-0.5	.28	.49	

WATER FEATURES						SOIL FEATURES		
I HVUIDIOUIC I	Depth to seasonal high water table (Feet)	Flooding		Ponding		Hydric		
		Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)	
Amenia	С	1.5-3.0	None		None		No	
Georgia	С	1.5-3.0	None		None		No	

	LAND USE LIMITA	AGRICULTURAL YIELD DATA			
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Georgia	Dwellings with basements:	Very limited	Depth to saturated zone	Pasture	8.5 AUM
Amenia	Dwellings with basements:	Very limited	Depth to saturated zone	Grass-legume hay	4.5 Tons
Georgia	ŭ	Very limited	Slope	Corn silage	22 Tons
· ·	Pond reservoir areas:	,	'	Alfalfa hay	4.5 Tons
Amenia	Pond reservoir areas:	Very limited	Slope	Corn silage	22 Tons
				Alfalfa hay	4.5 Tons
				Grass hay	4 Tons
				Grass-clover	6.4 AUM
				Grass-legume hay	4 Tons

	Management		WOODLAND MANA	AGEMENT .
Soil name	concern	Rating	Reason	Vermont natural communities
Georgia	Harvest equip operability:	Well suited		Mesic Maple-Ash-Hickory-Oak Forest,
Amenia	Harvest equip operability:	Well suited		Rich Northern Hardwood Forest, Sugar Maple-White Ash Northern Hardwood



Soil Fact Sheet - Continued

Rutland County, Vermont

Georgia Road suitability: Moderately suited Slope Forest

Amenia Road suitability: Moderately suited Slope

Georgia Erosion hazard (off-road): Slight Amenia Erosion hazard (off-road): Slight

Distribution Generation Date: 1/28/2015