

97C: Warwick-Quonset complex, 8 to 15 percent slopes

The Quonset component makes up 40 percent of the map unit. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. This component is on knolls on terraces on river valleys. The parent material consists of sandy and gravelly glaciofluvial deposits. Depth to a root restrictive layer is greater than 60 inches.

The Warwick component makes up 40 percent of the map unit. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is high. This component is on knolls on terraces on river valleys. The parent material consists of coarse-loamy glaciofluvial deposits. Depth to a root restrictive layer is greater than 60 inches.

Important farmland classification: Sta	atewide <u>Land capabilit</u>	<u>/:</u> 4 s <u>V</u>	/ermont Agricultural Value Group: 7
--	-------------------------------	------------------------	-------------------------------------

Vermont Residential Onsite Waste Disposal Group and Subgroup: la

This unit is well suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The rapid permeability in the substratum is a concern. Backfilling absorption trenches with at least one foot of finer textured material or other site modifications may be necessary to slow the percolation rate enough to allow for thorough filtering of effluent.

PHYSICAL and CHEMICAL PROPERTIES EROSION FACTORS									
Soil name	Depth	Typical	Clay	Soil	Permeability (In/Hr)	Organic matter	<u>LROSION I ACTORS</u>		
Soli Hame	(ln)	texture	(Pcť)	, i leaction		(Pct)	Kw	Kf	Т
Quonset	0-7	GRV-LS	2-7	4.5 - 5.5	2-20	1.0-7.0	.05	.15	5
	7-20	CNV-LS	1-4	4.5 - 6.0	2-20	0.5-2.0	.05	.17	
	20-60	CNV-COS	0-2	5.1 - 6.5	20-100	0.0-0.5	.02	.02	
Warwick	0-6	CN-FSL	3-8	4.5 - 6.5	2-6	1.0-5.0	.15	.24	3
	6-27	CNV-FSL	3-8	4.5 - 6.0	2-6	0.5-2.0	.10	.32	
	27-60	CNV-S	0-3	4.5 - 6.5	20-100	0.0-0.5	.02	.02	

	WATER FEATURES					SOIL FEATURES		
	Hydrologic	Depth to seasonal	Floo	ding	Pon	ding	Hydric	
Soil name	group	high water table (Feet)	Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)
Quonset	Α		None		None		No	
Warwick	Α		None		None		No	

	LAND USE LIMITA	AGRICULTURAL Y	AGRICULTURAL YIELD DATA		
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Warwick	Dwellings with basements:	Somewhat limited	Slope	Grass-clover	2.5 AUM
Quonset	Dwellings with basements:	Somewhat limited	Slope	Grass hay	3 Tons
Warwick	Pond reservoir areas:	Very limited	Seepage	Alfalfa hay	3.5 Tons
		,		Grass-legume hay	3 Tons
Quonset	Pond reservoir areas:	Very limited	Seepage	Grass-clover	4.8 AUM

	Management	<u>w</u>	OODLAND MANA	GEMENT
Soil name	concern	Rating	Reason	Vermont natural communities
Warwick	Harvest equip operability:	Well suited		White Pine-Red Oak-Black Oak Forest,
Quonset	Harvest equip operability:	Well suited		White Pine-Northern Hardwood Forest Variant
Narwick	Road suitability:	Moderately suited	Slope	
Quonset	Road suitability:	Moderately suited	Slope	
Varwick	Erosion hazard (off-road):	Slight		
Quonset	Erosion hazard (off-road):	Slight		