

97D: Warwick-Quonset complex, 15 to 25 percent slopes

The Quonset component makes up 40 percent of the map unit. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. This component is on terraces on river valleys. The parent material consists of sandy and gravelly glaciofluvial deposits. Depth to a root restrictive layer is greater than 60 inches.

The Warwick component makes up 40 percent of the map unit. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is high. This component is on terraces on river valleys. The parent material consists of coarse-loamy glaciofluvial deposits. Depth to a root restrictive layer is greater than 60 inches.

Important farmland classification: NPSL

Land capability: 6 s

Vermont Agricultural Value Group: 8

Vermont Residential Onsite Waste Disposal Group and Subgroup: 1b

This unit is well suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The rapid permeability in the substratum and slopes greater than 20 percent in some areas are concerns. Backfilling absorption trenches with at least one foot of finer textured material or other site modifications may be necessary to slow the percolation rate enough to allow for thorough filtering of effluent. There may be less-sloping areas within the unit that are suitable for siting a septic system, or, if feasible, cut and fill site modifications may produce an acceptable area within the unit. An erosion prevention and sediment control plan is required by the State for construction on sites over 20 percent slope.

PHYSICAL and CHEMICAL PROPERTIES							EROSION FACTORS		
Soil name	Depth (In)	Typical texture	Clay (Pct)	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	Kw	Kf	T
Quonset	0-7	GRV-LS	2-7	4.5 - 5.5	2-20	1.0-7.0	.05	.15	5
	7-20	CNV-LS	1-4	4.5 - 6.0	2-20	0.5-2.0	.05	.17	
	20-60	CNV-COS	0-2	5.1 - 6.5	20-100	0.0-0.5	.02	.02	
Warwick	0-6	CN-FSL	3-8	4.5 - 6.5	2-6	1.0-5.0	.15	.24	3
	6-27	CNV-FSL	3-8	4.5 - 6.0	2-6	0.5-2.0	.10	.32	
	27-60	CNV-S	0-3	4.5 - 6.5	20-100	0.0-0.5	.02	.02	

WATER FEATURES						SOIL FEATURES		
Soil name	Hydrologic group	Depth to seasonal high water table (Feet)	Flooding		Ponding		Hydric soil?	Depth to bedrock (range in inches)
			Frequency	Duration	Frequency	Duration		
Quonset	A	---	None		None		No	---
Warwick	A	---	None		None		No	---

LAND USE LIMITATIONS				AGRICULTURAL YIELD DATA	
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Warwick	Dwellings with basements:	Very limited	Slope	Grass-clover	2 AUM
Quonset	Dwellings with basements:	Very limited	Slope	Grass-legume hay	2.5 Tons
Warwick	Pond reservoir areas:	Very limited	Seepage	Grass-clover	4 AUM
Quonset	Pond reservoir areas:	Very limited	Seepage	Grass hay	2.5 Tons
				Alfalfa hay	3 Tons

WOODLAND MANAGEMENT				
Soil name	Management concern	Rating	Reason	Vermont natural communities
Warwick	Harvest equip operability:	Moderately suited	Slope	White Pine-Red Oak-Black Oak Forest, White Pine-Northern Hardwood Forest Variant
Quonset	Harvest equip operability:	Moderately suited	Slope	
Warwick	Road suitability:	Poorly suited	Slope	
Quonset	Road suitability:	Poorly suited	Slope	
Warwick	Erosion hazard (off-road):	Moderate	Slope/erodibility	
Quonset	Erosion hazard (off-road):	Moderate	Slope/erodibility	