

92D: Buckland loam, 15 to 25 percent slopes

The Buckland component makes up 85 percent of the map unit. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately low. This component is on hills on glaciated uplands. The parent material consists of loamy lodgment till derived from mica schist and/or loamy lodgment till derived from limestone and/or loamy lodgment till derived from phyllite. Depth to a root restrictive layer, densic material, is 20 to 36 inches.

Important farmland classification: NPSL	Land capability: 4 e	Vermont Agricultural Value Group: 8
---	----------------------	-------------------------------------

Vermont Residential Onsite Waste Disposal Group and Subgroup: IIIe

This unit is marginally suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The depth to the seasonal high water table and slopes greater than 20 percent in some areas are the major limitations. A detailed, site-specific analysis is generally required. On-site groundwater level monitoring and determination of induced groundwater mounding is often necessary to establish the suitability of this unit. Curtain drains may help lower the water table to an acceptable level. There may be less-sloping areas within the unit that are suitable for siting a septic system, or, if feasible, cut and fill site modifications may produce an acceptable area within the unit. An erosion prevention and sediment control plan is required by the State for construction on sites over 20 percent slope.

PHYSICAL and CHEMICAL PROPERTIES							EDOS	EROSION FACTORS	
Soil name	Depth	Typical	Clay	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	<u>EROSION FACTORS</u>		
	(ln)		(Pct)				Kw	Kf	Т
Buckland	0-8	L	2-20	5.6 - 7.3	0.1-14	2.0-8.0	.32	.32	3
	8-14	FSL	2-20	5.6 - 7.3	0.1-14	0.5-3.0	.55	.55	
	14-21	SIL	2-20	5.6 - 7.3	0.1-14	0.2-2.0	.64	.64	
	21-25	FSL	2-20	5.6 - 7.3	0.1-14	0.2-2.0	.64	.64	
	25-65	L	2-20	5.6 - 7.3	0.01-1	0.0-0.3	.55	.55	

WATER FEATURES						SOIL FEATURES		
Soil name	i i ivalolodic i	Hydrologic group Depth to seasonal high water table (Feet)	Flooding		Ponding		Hydric	
	group		Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)
Buckland	C/D	1.0-2.0	None		None		No	

	LAND USE LIMITAT	AGRICULTURAL YIELD DATA			
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Buckland	Dwellings with basements:	Very limited	Slope	Corn silage	18 Tons
Buckland	Pond reservoir areas:	voir areas: Very limited	Slope	Grass-legume hay	2.5 Tons
Tota reservoir areas.	Tond reservoir areas.			Grass-clover	4 AUM

	Management	<u>v</u>	VOODLAND MANAGEMEN	<u>ır</u>
Soil name	concern	Rating	Reason	Vermont natural communities
Buckland	Harvest equip operability:	Moderately suited	30-60cm to water table for the second	Northern Hardwood Forest, Rich Northern Hardwood Forest,
Buckland	Road suitability:	Poorly suited	Slope	Sugar Maple-White Ash Northern Hardwood Forest Variant
Buckland	Erosion hazard (off-road):	Moderate	Slope/erodibility	Troicst valiant