

68D: Taconic-Hubbardton-Rock outcrop complex, 8 to 25 percent slopes

The Taconic component makes up 50 percent of the map unit. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is low. This component is on hills on glaciated uplands, mountains on glaciated uplands. The parent material consists of coarse-loamy till. Depth to a root restrictive layer, bedrock, lithic, is 10 to 20 inches.

The Hubbardton component makes up 20 percent of the map unit. The natural drainage class is excessively drained. Water movement in the most restrictive layer is low. This component is on hills on glaciated uplands, mountains on glaciated uplands. The parent material consists of coarse-loamy till. Depth to a root restrictive layer, bedrock, lithic, is 2 to 10 inches.

The Rock outcrop component makes up 20 percent of the map unit. Water movement in the most restrictive layer is low. This component is on hills on glaciated uplands, mountains on glaciated uplands. Depth to a root restrictive layer0 inches, bedrock, lithic..

Important farmland classification: N	IPSL	Land capability: 7 e	е	Vermont Agricultural Value Group:	10	l
--------------------------------------	------	----------------------	---	-----------------------------------	----	---

Vermont Residential Onsite Waste Disposal Group and Subgroup: IVc

This unit is generally not suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The very shallow to shallow depth to bedrock is the limiting condition.

PHYSICAL and CHEMICAL PROPERTIES								EROSION FACTORS	
Soil name	Depth	Typical	Clay	Soil	Soil Permeability eaction (In/Hr)	Organic matter (Pct)	<u>EROSION FACTORS</u>		
30ii Hame	(ln)	texture	(Pcť)				Kw	Kf	Т
Taconic	0-2	CN-L	10-27	4.5 - 5.5	0.6-6	2.0-6.0	.15	.28	1
	2-19	CNV-L	10-27	4.5 - 5.5	0.6-6	0.0-1.0	.15	.49	
	19-29	UWB			0.01-20				
Hubbardton	0-2	CNV-SIL	3-18	4.5 - 5.5	0.6-6	1.0-4.0	.20	.49	1
	2-6	CNV-SIL	3-18	4.5 - 5.5	0.6-6	0.0-1.0	.20	.55	
	6-16	UWB			0.01-20				
Rock outcrop	0-60	UWB			0.01-20				

WATER FEATURES							SOIL FEATURES		
	Hydrologic	Depth to seasonal	Floo	ding	Pon	ding	Hydric		
Soil name	group	high water table (Feet)	Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)	
Taconic	D		None		None		No	10-20	
Hubbardton	D		None		None		No	2-10	
Rock outcrop			None				Unranked	0	

	LAND USE LIMITA	AGRICULTURAL YIELD DATA			
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Hubbardton	Dwellings with basements:	Very limited	Depth to hard bedrock	Pasture	1.5 AUM
Taconic	Dwellings with basements:	Very limited	Depth to hard bedrock		
Hubbardton	Pond reservoir areas:	Very limited	Slope		
Taconic	Pond reservoir areas:	Very limited	Slope		

	Management		WOODLAND MANAGE	<u>EMENT</u>
Soil name	concern	Rating	Reason	Vermont natural communities
Hubbardton	Harvest equip operability:	Well suited		Dry Oak-Hickory-Hophornbeam Forest,
Taconic	Harvest equip operability:	Well suited		Mesic Red Oak-Northern Hardwood Forest, Dry Oak Forest,
Hubbardton	Road suitability:	Poorly suited	Slope	Red Cedar Woodland,
Taconic	Road suitability:	Poorly suited	Slope	Mesic Maple-Ash-Hickory-Oak Forest,
Hubbardton	Erosion hazard (off-road):	Moderate	Slope/erodibility	Temperate Acidic Outcrop, Temperate Acidic Cliff,



Soil Fact Sheet - Continued

Windham County, Vermont

Taconic

Erosion hazard (off-road): Moderate

Slope/erodibility

Dry Oak Woodland