

1C: Hitchcock silt loam, 8 to 15 percent slopes

The Hitchcock component makes up 85 percent of the map unit. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately low. This component is on strongly sloping lake terraces on lake plains. The parent material consists of silty glaciolacustrine deposits. Depth to a root restrictive layer is greater than 60 inches.

| Important farmland classification: Statewide | Land capability: 3 e | Vermont Agricultural Value Group: 5 |
|--|----------------------|-------------------------------------|
|--|----------------------|-------------------------------------|

Vermont Residential Onsite Waste Disposal Group and Subgroup: Ila

This unit is moderately suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The slow permeability in the substratum is the primary concern. Mound system construction and other site modifications may be necessary.

| PHYSICAL and CHEMICAL PROPERTIES | | | | | | | EROSION FACTORS | | |
|----------------------------------|-------|--------------|---------------|--------------------------|-------------------------|----------------------------|-------------------------|-----|---|
| Soil name | Depth | Typical | Clay (Pct) | Soil reaction (pH) | Permeability (In/Hr) | Organic matter (Pct) | <u>EROSION I ACTORS</u> | | |
| | (In) | (In) texture | | | | | Kw | Kf | Т |
| Hitchcock | 0-9 | SIL | 3-10 | 4.5 - 6.5 | 0.6-2 | 1.0-5.0 | .49 | .49 | 5 |
| | 9-26 | SIL | 3-15 | 4.5 - 6.5 | 0.6-2 | 0.5-3.0 | .55 | .55 | |
| | 26-65 | SIL | 3-15 | 4.5 - 6.5 | 0.06-0.6 | 0.0-1.0 | .64 | .64 | |

| WATER FEATURES | | | | | | | SOIL FEATURES | | |
|----------------|------------|------------------------------|-----------|----------|-----------|----------|---------------|------------------------------------|--|
| | Hydrologic | Hydrologic Depth to seasonal | | ding Por | | Ponding | | | |
| Soil name | group | high water table (Feet) | Frequency | Duration | Frequency | Duration | soil? | Depth to bedrock (range in inches) | |
| Hitchcock | С | | None | | None | | No | | |

| • | LAND USE LIMITAT | AGRICULTURAL YIE | AGRICULTURAL YIELD DATA | | |
|-----------|---------------------------|------------------|-------------------------|------------------|--------------|
| Soil name | Land use | Rating | Reason ** | Crop name | Yield / acre |
| Hitchcock | Dwellings with basements: | Somewhat limited | Slope | Pasture | 7.5 AUM |
| Hitchcock | Pond reservoir areas: | Very limited | Slope | Grass-legume hay | 4 Tons |
| THEHOOCK | Fond reservoir areas. | very minica | Оюрс | Grass hay | 4.5 Tons |
| | | | | Corn silage | 22 Tons |
| | | | | Alfalfa hay | 4 Tons |

| | Management | <u>v</u> | VOODLAND MANA | <u>GEMENT</u> |
|-----------|----------------------------|-------------------|-------------------|--|
| Soil name | concern | Rating | Reason | Vermont natural communities |
| Hitchcock | Harvest equip operability: | Moderately suited | Low strength | Northern Hardwood Forest, |
| Hitchcock | Road suitability: | Moderately suited | Slope | Mesic Maple-Ash-Hickory-Oak Forest, Rich Northern Hardwood Forest |
| Hitchcock | Erosion hazard (off-road): | Moderate | Slope/erodibility | Titili Worthern Hardwood Forest |