

25B: Buckland loam, 3 to 8 percent slopes

The Buckland component makes up 84 percent of the map unit. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately low. This component is on hills on glaciated uplands. The parent material consists of loamy lodgment till derived from mica schist and/or loamy lodgment till derived from limestone and/or loamy lodgment till derived from phyllite. Depth to a root restrictive layer, densic material, is 20 to 40 inches.

Important farmland classification: Prime

Land capability: 2 w

Vermont Agricultural Value Group: 3

Vermont Residential Onsite Waste Disposal Group and Subgroup: IIIc

This unit is marginally suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The depth to the seasonal high water table in association with the minimal slope is the major limitation. A detailed, site-specific analysis is generally required. On-site groundwater level monitoring and determination of induced groundwater mounding is often necessary to establish the suitability of this unit. Curtain drains may help lower the water table to an acceptable level, however, the minimal slope may prevent their use in many areas.

PHYSICAL and CHEMICAL PROPERTIES							EROSION FACTORS		
Soil name	Depth (In)	Typical texture	Clay (Pct)	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	Kw	Kf	T
Buckland	0-8	L	2-20	5.6 - 7.3	0.1-14	2.0-8.0	.32	.32	3
	8-14	FSL	2-20	5.6 - 7.3	0.1-14	0.5-3.0	.55	.55	
	14-21	SIL	2-20	5.6 - 7.3	0.1-14	0.2-2.0	.64	.64	
	21-25	FSL	2-20	5.6 - 7.3	0.1-14	0.2-2.0	.64	.64	
	25-65	L	2-20	5.6 - 7.3	0.01-1	0.0-0.3	.55	.55	

WATER FEATURES						SOIL FEATURES		
Soil name	Hydrologic group	Depth to seasonal high water table (Feet)	Flooding		Ponding		Hydric soil?	Depth to bedrock (range in inches)
			Frequency	Duration	Frequency	Duration		
Buckland	C/D	1.0-2.0	None		None		No	---

LAND USE LIMITATIONS				AGRICULTURAL YIELD DATA	
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Buckland	Dwellings with basements:	Very limited	Depth to saturated zone	Grass-clover	5.6 AUM
Buckland	Pond reservoir areas:	Somewhat limited	Seepage	Grass-legume hay	3.5 Tons
				Corn silage	22 Tons

WOODLAND MANAGEMENT				
Soil name	Management concern	Rating	Reason	Vermont natural communities
Buckland	Harvest equip operability:	Moderately suited	30-60cm to water table	Northern Hardwood Forest, Rich Northern Hardwood Forest, Sugar Maple-White Ash Northern Hardwood Forest Variant
Buckland	Road suitability:	Moderately suited	Wetness	
Buckland	Erosion hazard (off-road):	Slight		