

54C: Tunbridge-Lyman complex, 8 to 15 percent slopes, rocky

The Tunbridge, rocky component makes up 50 percent of the map unit. The natural drainage class is well drained. Water movement in the most restrictive layer is low. This component is on mountains on glaciated uplands, hills on glaciated uplands. The parent material consists of loamy supraglacial till derived from granite and gneiss and/or loamy supraglacial till derived from phyllite and/or loamy supraglacial till derived from mica schist. Depth to a root restrictive layer, bedrock, lithic, is 20 to 40 inches.

The Lyman, rocky component makes up 33 percent of the map unit. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is low. This component is on hills on glaciated uplands, mountains on glaciated uplands. The parent material consists of loamy supraglacial till derived from granite and gneiss and/or loamy supraglacial till derived from mica schist. Depth to a root restrictive layer, bedrock, lithic, is 11 to 24 inches.

Important farmland classification:	Statewide	Land capability: 3 e	Vermont Agricultural Value Group: 5
------------------------------------	-----------	----------------------	-------------------------------------

Vermont Residential Onsite Waste Disposal Group and Subgroup: Ilc

This unit is moderately suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The depth to bedrock in some areas is the primary concern. A significant percentage of this map unit has sufficient soil depth over bedrock to accept a range of designs. On-site investigations can help avoid areas with limited depth to bedrock. Additional fill material may be needed in some areas in order to meet the separation distance requirement between the bottom of the leachfield and bedrock.

PHYSICAL and CHEMICAL PROPERTIES								EDOCION EACTORS	
Soil name	Depth	Typical	Clay (Pct)	Soil reaction (pH)	Permeability (In/Hr)	Organic matter (Pct)	EROSION FACTORS		
30ii Hairie	(ln)	texture					Kw	Kf	Т
Tunbridge, rocky	0-3	MPM		3.5 - 5.5	1-14	35-95			2
	3-5	HPM		3.5 - 5.5	1-14	35-95			
	5-8	FSL	1-10	3.5 - 6.0	0.1-14	1.4-5.7	.37	.37	
	8-11	FSL	1-10	3.5 - 6.0	0.1-14	3.1-25	.32	.32	
	11-26	FSL	1-10	3.5 - 6.0	0.1-14	2.2-18	.37	.37	
	26-28	FSL	1-10	5.1 - 6.5	0.1-14	1.0-4.2	.43	.43	
	28-38	BR			0.001-14				
man, rocky	0-1	MPM		3.5 - 5.5	1-14	35-95			1
	1-3	L	1-10	3.5 - 6.0	0.1-14	4.0-20	.32	.32	
	3-5	FSL	1-10	3.5 - 6.0	0.1-14	1.4-5.7	.37	.37	
	5-7	L	1-10	3.5 - 6.0	0.1-14	3.1-25	.32	.32	
	7-11	L	1-10	3.5 - 6.0	0.1-14	2.2-18	.32	.32	
	11-18	CN-L	1-10	3.5 - 6.0	0.1-14	2.2-7.0	.24	.32	
	18-28	BR			0.001-14				

WATER FEATURES						SOIL FEATURES		
	Hydrologic	Depth to seasonal	Flooding		Ponding		Hydric	
Soil name			Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)
Tunbridge, rocky	С		None		None		No	20-40
Lyman, rocky	D		None		None		No	11-24

	LAND USE LIMITA	AGRICULTURAL YIELD DATA			
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Tunbridge, rocky	Dwellings with basements:	Very limited	Depth to hard bedrock	Grass-clover	5.6 AUM
Lyman, rocky	Dwellings with basements:	Very limited	Depth to hard bedrock	Corn silage	18 Tons
Tunbridge, rocky	Pond reservoir areas:	Very limited	Slope	Grass-legume hay	3.5 Tons
Lyman, rocky	Pond reservoir areas:	Very limited Slope	Grass-clover	4.8 AUM	
Lyman, rocky	Folia reservoir areas.	very innited	Оюрс	Corn silage	12 Tons
				Grass-legume hay	2.5 Tons

Soil Fact Sheet - Continued

Windsor County, Vermont

	Management	<u>v</u>	VOODLAND MANAGE	<u>EMENT</u>
Soil name	concern	Rating	Reason	Vermont natural communities
Tunbridge	Harvest equip operability:	Well suited		Northern Hardwood Forest,
Lyman	Harvest equip operability:	Well suited		Hemlock-Northern Hardwood Forest, Mesic Red Oak-Northern Hardwood Forest,
Tunbridge	Road suitability:	Moderately suited	Slope	Beech-Red Maple-Hemlock-Northern Hardwood
Lyman	Road suitability:	Moderately suited	Slope	Forest Variant,
Tunbridge	Erosion hazard (off-road):	Slight		Hemlock Forest
Lyman	Erosion hazard (off-road):	Moderate	Slope/erodibility	