

80C: Macomber-Taconic complex, 8 to 15 percent slopes, very rocky

The Macomber, very rocky component makes up 55 percent of the map unit. The natural drainage class is well drained. Water movement in the most restrictive layer is low. This component is on strongly sloping hills on glaciated uplands. The parent material consists of loamy till. Depth to a root restrictive layer, bedrock, lithic, is 20 to 40 inches.

The Taconic, very rocky component makes up 30 percent of the map unit. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is low. This component is on strongly sloping hills on glaciated uplands. The parent material consists of loamy till. Depth to a root restrictive layer, bedrock, lithic, is 10 to 20 inches.

Important farmland classification: NPSL	Land capability: 6 s	Vermont Agricultural Value Group: 10
---	----------------------	--------------------------------------

Vermont Residential Onsite Waste Disposal Group and Subgroup: IIc

This unit is moderately suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The depth to bedrock in some areas is the primary concern. A significant percentage of this map unit has sufficient soil depth over bedrock to accept a range of designs. On-site investigations can help avoid areas with limited depth to bedrock. Additional fill material may be needed in some areas in order to meet the separation distance requirement between the bottom of the leachfield and bedrock.

PHYSICAL and CHEMICAL PROPERTIES							EDOCION EACTORS		
Callmana	Depth	Typical	Clay	Soil	Permeability (In/Hr)	Organic matter (Pct)	EROSION FACTORS		
Soil name	(In)	texture	(Pct)	reaction (pH)			Kw	Kf	Т
Macomber, very rocky	0-3	MPM		3.2 - 5.7	2-6	25-100			2
	3-7	SIL	10-27	4.5 - 5.5	0.6-2	2.0-6.0	.28	.28	
	7-33	CNV-SIL	10-27	4.5 - 5.5	0.6-2	0.0-1.0	.15	.49	
	33-43	UWB			0.01-20				
Taconic, very rocky	0-3	MPM		3.2 - 5.7	2-6	25-100			1
	3-6	CN-SIL	10-27	4.5 - 5.5	0.6-6	2.0-6.0	.20	.37	
	6-15	CNV-SIL	10-27	4.5 - 5.5	0.6-6	0.0-1.0	.20	.49	
	15-25	UWB			0.01-20				

		<u>WATE</u>	R FEATURES				SOIL	FEATURES
	Hydrologic	Depth to seasonal	Floo	ding	Pon	ding	Hydric	
Soil name	group	high water table (Feet)	Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)
Macomber, very rocky	С		None		None		No	20-40
Taconic, very rocky	D		None		None		No	10-20

	LAND USE LIMITA	<u>TIONS</u>		AGRICULTURA	L YIELD DATA
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Macomber, very	Dwellings with basements:	Very limited	Depth to hard bedrock	Pasture	3.4 AUM
rocky		•		Pasture	2.4 AUM
Taconic, very rocky	Dwellings with basements:	Very limited	Depth to hard bedrock	•	
Macomber, very rocky	Pond reservoir areas:	Very limited	Slope		
Taconic, very rocky	Pond reservoir areas:	Very limited	Slope		

	Management	<u>v</u>	OODLAND MANA	GEMENT
Soil name	concern	Rating	Reason	Vermont natural communities
Macomber	Harvest equip operability:	Well suited		Northern Hardwood Forest,
Taconic	Harvest equip operability:	Well suited		Mesic Red Oak-Northern Hardwood Forest, Dry Oak-Hickory-Hophornbeam Forest,
Macomber	Road suitability:	Moderately suited	Slope	Red Cedar Woodland,
aconic	Road suitability:	Moderately suited	Slope	Mesic Maple-Ash-Hickory-Oak Forest,
//acomber	Erosion hazard (off-road):	Slight		Temperate Acidic Outcrop, Temperate Acidic Cliff
aconic	Erosion hazard (off-road):	Slight		

