

81D: Taconic-Hubbardton-Rock outcrop complex, 8 to 25 percent slopes

The Taconic component makes up 50 percent of the map unit. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is low. This component is on strongly sloping to moderately steep hills on glaciated uplands. The parent material consists of loamy till. Depth to a root restrictive layer, bedrock, lithic, is 10 to 20 inches.

The Hubbardton component makes up 20 percent of the map unit. The natural drainage class is excessively drained. Water movement in the most restrictive layer is low. This component is on strongly sloping to moderately steep hills on glaciated uplands. The parent material consists of loamy till. Depth to a root restrictive layer, bedrock, lithic, is 2 to 10 inches.

The Rock outcrop component makes up 15 percent of the map unit. Water movement in the most restrictive layer is low. This component is on strongly sloping to moderately steep hills. Depth to a root restrictive layer0 inches, bedrock, lithic,.

Important farmland classification: N	NPSL	Land capability: 7 s	Vermont Agricultural Value Group: 10

Vermont Residential Onsite Waste Disposal Group and Subgroup: IVc

This unit is generally not suited as a site for soil-based residential wastewater disposal systems, based on a review by the Natural Resources Conservation Service of criteria set forth in the Vermont 2007 Environmental Protection Rules. The very shallow to shallow depth to bedrock is the limiting condition.

PHYSICAL and CHEMICAL PROPERTIES							EDOCION EACTORS		
0.1	Depth	Typical	Clay	Soil	Permeability (In/Hr)	Organic matter (Pct)	EROSION FACTORS		
Soil name	(In)	texture	(Pct)	reaction (pH)			Kw	Kf	Т
Taconic	0-3	MPM		3.2 - 5.7	2-6	25-100			1
	3-6	CN-SIL	10-27	4.5 - 5.5	0.6-6	2.0-6.0	.20	.37	
	6-15	CNV-SIL	10-27	4.5 - 5.5	0.6-6	0.0-1.0	.20	.49	
	15-25	UWB			0.01-20				
Hubbardton	0-3	MPM		3.2 - 5.7	2-6	25-100			1
	3-6	CNV-SIL	3-18	4.5 - 5.5	0.6-6	0.0-1.0	.20	.55	
	6-16	UWB			0.01-20				
Rock outcrop	0-65	UWB			0.01-20				

	WATER FEATURES					SOIL FEATURES			
	Hydrologic	Depth to seasonal	Flooding		Ponding		Hydric		
Soil name	group	high water table (Feet)	Frequency	Duration	Frequency	Duration	soil?	Depth to bedrock (range in inches)	
Taconic	D		None		None		No	10-20	
Hubbardton	D		None		None		No	2-10	
Rock outcrop			None		None		Unranked	0	

	LAND USE LIMITA	AGRICULTURAL YIELD DATA			
Soil name	Land use	Rating	Reason **	Crop name	Yield / acre
Taconic Hubbardton	Dwellings with basements: Dwellings with basements:	,	Depth to hard bedrock Depth to hard bedrock	Pasture	1.5 AUM
Taconic Hubbardton	Pond reservoir areas: Pond reservoir areas:	Very limited Very limited	Slope Slope		

	Management		WOODLAND MANA	GEMENT
Soil name	concern	Rating	Reason	Vermont natural communities
Taconic	Harvest equip operability:	Well suited		Dry Oak-Hickory-Hophornbeam Forest,
Hubbardton	Harvest equip operability:	Well suited		Mesic Red Oak-Northern Hardwood Forest, Dry Oak Forest,
Taconic	Road suitability:	Poorly suited	Slope	Red Cedar Woodland,
Hubbardton	Road suitability:	Poorly suited	Slope	Mesic Maple-Ash-Hickory-Oak Forest,



Soil Fact Sheet - Continued

Windsor County, Vermont

Taconic Erosion hazard (off-road): Moderate
Hubbardton Erosion hazard (off-road): Moderate

Slope/erodibility Slope/erodibility Temperate Acidic Outcrop, Temperate Acidic Cliff, Dry Oak Woodland